

Novel Imidazolidinones as NS3-Serine Protease Inhibitors of Hepatitis C Virus

Field of Invention

The present invention relates to novel hepatitis C virus ("HCV") protease inhibitors, pharmaceutical compositions containing one or more such inhibitors, methods of preparing such inhibitors and methods of using such inhibitors to treat hepatitis C and related disorders. This invention specifically discloses novel peptide compounds as inhibitors of the HCV NS3/NS4a serine protease.

Background of the Invention

Hepatitis C virus (HCV) is a (+)-sense single-stranded RNA virus that has been implicated as the major causative agent in non-A, non-B hepatitis (NANBH), particularly in blood-associated NANBH (BB-NANBH)(see, International Patent Application Publication No. WO 89/04669 and European Patent Application Publication No. EP 381 216). NANBH is to be distinguished from other types of viral-induced liver disease, such as hepatitis A virus (HAV), hepatitis B virus (HBV), delta hepatitis virus (HDV), cytomegalovirus (CMV) and Epstein-Barr virus (EBV), as well as from other forms of liver disease such as alcoholism and primary biliar cirrhosis.

Recently, an HCV protease necessary for polypeptide processing and viral replication has been identified, cloned and expressed; (see, e.g., U.S. Patent No. 5,712,145). This approximately 3000 amino acid polyprotein contains, from the amino terminus to the carboxy terminus, a nucleocapsid protein (C), envelope proteins (E1 and E2) and several non-structural proteins (NS1, 2, 3, 4a, 5a and 5b). NS3 is an approximately 68 kda protein, encoded by approximately 1893 nucleotides of the HCV genome, and has two distinct domains: (a) a serine protease domain consisting of approximately 200 of the N-terminal amino acids; and (b) an RNA-dependent ATPase domain at the C-terminus of the protein. The

NS3 protease is considered a member of the chymotrypsin family because of similarities in protein sequence, overall three-dimensional structure and mechanism of catalysis. Other chymotrypsin-like enzymes are elastase, factor Xa, thrombin, trypsin, plasmin,

urokinase, tPA and PSA. The HCV NS3 serine protease is responsible for proteolysis of the polypeptide (polyprotein) at the NS3/NS4a, NS4a/NS4b, NS4b/NS5a and NS5a/NS5b junctions and is thus responsible for generating four viral proteins during viral replication. This has made the HCV NS3 serine protease an attractive target for antiviral chemotherapy.

It has been determined that the NS4a protein, an approximately 6 kda polypeptide, is a co-factor for the serine protease activity of NS3. Autocleavage of the NS3/NS4a junction by the NS3/NS4a serine protease occurs intramolecularly (*i.e.*, *cis*) while the other cleavage sites are processed intermolecularly (*i.e.*, *trans*).

Analysis of the natural cleavage sites for HCV protease revealed the presence of cysteine at P1 and serine at P1' and that these residues are strictly conserved in the NS4a/NS4b, NS4b/NS5a and NS5a/NS5b junctions. The NS3/NS4a junction contains a threonine at P1 and a serine at P1'. The Cys→Thr substitution at NS3/NS4a is postulated to account for the requirement of *cis* rather than *trans* processing at this junction. See, *e.g.*, Pizzi *et al.* (1994) *Proc. Natl. Acad. Sci. (USA)* **91**:888-892, Failla *et al.* (1996) *Folding & Design* **1**:35-42. The NS3/NS4a cleavage site is also more tolerant of mutagenesis than the other sites. See, *e.g.*, Kollykhalov *et al.* (1994) *J. Virol.* **68**:7525-7533. It has also been found that acidic residues in the region upstream of the cleavage site are required for efficient cleavage. See, *e.g.*, Komoda *et al.* (1994) *J. Virol.* **68**:7351-7357.


Inhibitors of HCV protease that have been reported include antioxidants (see, International Patent Application Publication No. WO 98/14181), certain peptides and peptide analogs (see, International Patent Application Publication No. WO 98/17679, Landro *et al.* (1997) *Biochem.* **36**:9340-9348, Ingallinella *et al.* (1998) *Biochem.* **37**:8906-8914, Llinàs-Brunet *et al.* (1998) *Bioorg. Med. Chem. Lett.* **8**:1713-1718), inhibitors based on the 70-amino acid polypeptide eglin c (Martin *et al.* (1998) *Biochem.* **37**:11459-11468, inhibitors affinity selected from

human pancreatic secretory trypsin inhibitor (hPSTI-C3) and minibody repertoires (MBip) (Dimasi *et al.* (1997) *J. Virol.* **71**:7461-7469), cV_HE2 (a "camelized" variable domain antibody fragment) (Martin *et al.* (1997) *Protein Eng.* **10**:607-614), and α 1-antichymotrypsin (ACT) (Elzouki *et al.* (1997) *J. Hepat.* **27**:42-28). A ribozyme designed to selectively destroy hepatitis C virus RNA has recently been disclosed (see, *BioWorld Today* **9**(217): 4 (November 10, 1998)).

Reference is also made to the PCT Publications, No. WO 98/17679, published April 30, 1998 (Vertex Pharmaceuticals Incorporated); WO 98/22496, published May 28, 1998 (F. Hoffmann-La Roche AG); and WO 99/07734, published February 18, 1999 (Boehringer Ingelheim Canada Ltd.).

HCV has been implicated in cirrhosis of the liver and in induction of hepatocellular carcinoma. The prognosis for patients suffering from HCV infection is currently poor. HCV infection is more difficult to treat than other forms of hepatitis due to the lack of immunity or remission associated with HCV infection. Current data indicates a less than 50% survival rate at four years post cirrhosis diagnosis. Patients diagnosed with localized resectable hepatocellular carcinoma have a five-year survival rate of 10-30%, whereas those with localized unresectable hepatocellular carcinoma have a five-year survival rate of less than 1%.


Reference is made to A. Marchetti *et al.*, *Synlett*, **S1**, 1000-1002 (1999) describing the synthesis of bicyclic analogs of an inhibitor of HCV NS3 protease. A compound disclosed therein has the formula:




Reference is also made to W. Han *et al.*, *Bioorganic & Medicinal Chem. Lett.* (2000) **10**, 711-713, which describes the preparation of certain α -

ketoamides, α -ketoesters and α -diketones containing allyl and ethyl functionalities.


Reference is also made to WO 00/09558 (Assignee: Boehringer Ingelheim Limited; Published February 24, 2000) which discloses peptide derivatives of the
5 formula:




where the various elements are defined therein. An illustrative compound of that series is:



Reference is also made to WO 00/09543 (Assignee: Boehringer Ingelheim Limited; Published February 24, 2000) which discloses peptide derivatives of the formula:



where the various elements are defined therein. An illustrative compound of that series is:



Current therapies for hepatitis C include interferon- α (INF $_{\alpha}$) and combination therapy with ribavirin and interferon. See, e.g., Beremguer *et al.* (1998) *Proc. Assoc. Am. Physicians* 110(2):98-112. These therapies suffer from a low sustained response rate and frequent side effects. See, e.g., Hoofnagle *et al.* (1997) *N. Engl. J. Med.* 336:347. Currently, no vaccine is available for HCV infection.

Pending and copending U. S. patent applications, Serial No. 60/220,109, filed July 21, 2000, Serial No. 60/220,107, filed July 21, 2000, Serial No. 60/220,108, filed July 21, 2000, Serial No. 60/220,101, filed July 21, 2000, Serial No. 60/254,869, filed December 12, 2000, Serial No. 60/194,607, filed April 5, 2000, and Serial No. 60/198,204, filed April 19, 2000 disclose various types of peptides as NS-3 serine protease inhibitors of hepatitis C virus.

There is a need for new treatments and therapies for HCV infection. It is, therefore, an object of this invention to provide compounds useful in the treatment or prevention or amelioration of one or more symptoms of hepatitis C.

It is a further object herein to provide methods of treatment or prevention or amelioration of one or more symptoms of hepatitis C.


A still further object of the present invention is to provide methods for modulating the activity of serine proteases, particularly the HCV NS3/NS4a serine protease, using the compounds provided herein.

Another object herein is to provide methods of modulating the processing of the HCV polypeptide using the compounds provided herein.

Summary of Invention

In its several embodiments, the present invention provides a novel class of inhibitors of the HCV protease, pharmaceutical compositions containing one or more of the compounds, methods of preparing pharmaceutical formulations comprising one or more such compounds, and methods of treatment, prevention or amelioration or one or more of the symptoms of hepatitis C. Also provided are methods of modulating the interaction of an HCV polypeptide with HCV protease. Among the compounds provided herein, compounds that inhibit HCV NS3/NS4a serine protease activity are preferred. The presently disclosed compounds generally contain about two or more amino acid residues and less than about twelve amino acid residues.

In its principal embodiment, the present invention provides an imidazolidinone of Formula I:



Formula I

wherein:

- 5 R^1 is COR^5 or $B(OR)_2$, wherein R^5 is selected from the group consisting of: H, OH, OR^8 , NR^9R^{10} , CF_3 , C_2F_5 , C_3F_7 , CF_2R^6 , R^6 , and COR^7 with R^7 being H, OH, OR^8 , CHR^9R^{10} , or NR^9R^{10} , wherein R^6 , R^8 , R^9 and R^{10} are independently selected from the group consisting of H, alkyl, aryl, heteroalkyl, heteroaryl, cycloalkyl, cycloalkyl, arylalkyl, heteroarylalkyl, $CH(R^1)COOR^{11}$, $CH(R^1)CONR^{12}R^{13}$,
 10 $CH(R^1)CONHCH(R^2)COO R^{11}$, $CH(R^1)CONHCH(R^2)CONR^{12}R^{13}$, $CH(R^1)CONHCH(R^2)R'$, $CH(R^1)CONHCH(R^2)CONHCH(R^3)COO R^{11}$, $CH(R^1)CONHCH(R^2)CONHCH(R^3)CONR^{12}R^{13}$, wherein R^1 , R^2 , R^3 , R' , R^{11} , R^{12} , and R^{13} are independently selected from a group consisting of H, alkyl, aryl, heteroalkyl, heteroaryl, cycloalkyl, alkyl-aryl, alkyl-heteroaryl, aryl-alkyl and heteroalkyl;
 15

Z is O, N, or CH;

X is selected from the group consisting of: C=O, C=S and $(CRR')_p$;

p is a number from 1-6;

G is selected from the group consisting of: H, alkyl, aryl, heteroalkyl, heteroaryl,

- 20 alkyl-aryl and alkyl-heteroaryl with the proviso that G may be additionally optionally and chemically-suitably substituted with U^{11} or U^{12} ;

R , R^2 , and R^3 may be the same or different and are independently selected from the group consisting of H; C1-C10 alkyl; C2-C10 alkenyl; C3-C8 cycloalkyl; C3-C8 heterocycloalkyl, alkoxy, aryloxy, alkylthio, arylthio, amino, amido, ester,

- 25 carboxylic acid, carbamate, urea, ketone, aldehyde, cyano, nitro;

(cycloalkyl)alkyl and (heterocycloalkyl)alkyl, wherein said cycloalkyl is made of three to eight carbon atoms, and zero to six oxygen, nitrogen, sulfur, or phosphorus atoms, and said alkyl is of one to six carbon atoms; aryl; heteroaryl; alkyl-aryl; and alkyl-heteroaryl;

- 5 R^4 maybe present or absent, and if R^4 is present, R^4 is selected from H, alkyl, aryl; and

Y is selected from the group consisting of H, alkyl, aryl, heteroalkyl, heteroaryl, cycloalkyl, arylalkyl, heteroarylalkyl, and U, where U is selected from alkyl-aryl, aryl-heteroaryl, alkyl-heteroaryl, alkylcarbonyl, arylalkylcarbonyl, arylcarbonyl, heteroalkylcarbonyl, heteroarylcarbonyl, cycloalkylcarbonyl, alkyloxycarbonyl, alkyl-aryloxycarbonyl, aryloxycarbonyl, arylalkyloxycarbonyl, heteroaryloxycarbonyl, heterocycloalkyloxycarbonyl, cycloalkyloxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, alkyl-arylaminocarbonyl, arylaminocarbonyl, arylalkylaminocarbonyl, heteroarylaminocarbonyl, cycloalkylaminocarbonyl, heterocycloalkylaminocarbonyl, alkylsulfonyl, arylsulfonyl, alkyl-arylsulfonyl, arylalkylsulfonyl, heteroarylsulfonyl, cycloalkylsulfonyl, and heterocycloalkylsulfonyl with the proviso that U may be additionally optionally and chemically-suitably substituted with U^{11} or U^{12} ; where

U^{11} is alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkyl-alkyl, heterocyclyl,

- 10 heterocyclylalkyl, aryl, alkylaryl, arylalkyl, heteroaryl, alkylheteroaryl, or heteroarylalkyl moiety, with the proviso that U^{11} may be additionally optionally substituted with U^{12} ; and

U^{12} is hydroxy, alkoxy, aryloxy, thio, alkylthio, arylthio, amino, alkylamino, arylamino, alkylsulfonyl, arylsulfonyl, alkylsulfonamido, arylsulfonamido, carboxy, carbalkoxy, carboxamido, alkoxycarbonylamino, alkoxycarbonyloxy, alkylureido, arylureido, halogen, cyano, or nitro moiety, with the proviso that the alkyl, alkoxy, and aryl may be additionally optionally substituted with moieties independently selected from U^{12} . Furthermore, it is to be generally understood that the alkyl, heteroalkyl, alkenyl, heteroalkenyl, aryl, heteroaryl, cycloalkyl and heterocycloalkyl


15
20
25
30

moieties may be optionally substituted, with the term "substituted" referring to optional and chemically-suitable substitution with one or more moieties selected


5

$$\begin{array}{c} \text{Y} \\ \diagdown \\ \text{R}^4 - \text{Z} - \text{C} - \text{R}^3 \\ \diagup \\ \text{R}^3 \end{array}$$


Some illustrative preferred examples for R^2 include the following:



Some illustrative preferred examples for G include the following:




Some illustrative preferred examples for R^3 include the following:



wherein R^{31} = H, alkyl or aryl.

In another embodiment, the present invention discloses imidazolidinones of

Formula II:




Formula II

wherein R^1 , R^2 , R^3 , R^4 , G, X, Y and Z are as defined above, t is a number from 1 to 3, and the P moiety or moieties (when t is 2 or 3) may be the same or different,

are amino acid residues represented by:



wherein V is selected from the group consisting of the following:



wherein R^{31} is H, alkyl or aryl.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which this invention belongs. Thus, for example, the term alkyl (including the alkyl portions of alkoxy) refers to a monovalent group derived from a straight or branched chain saturated hydrocarbon by the removal of a single atom having from 1 to 8 carbon atoms, preferably from 1 to 6;

aryl – represents a carbocyclic group having from 6 to 14 carbon atoms and having at least one benzenoid ring, with all available substitutable aromatic carbon atoms of the carbocyclic group being intended as possible points of attachment. Preferred aryl groups include phenyl, 1-naphthyl, 2-naphthyl and indanyl, and especially phenyl and substituted phenyl;

aralkyl – represents a moiety containing an aryl group linked via a lower alkyl;

alkylaryl – represents a moiety containing a lower alkyl linked via an aryl group;

cycloalkyl – represents a saturated carbocyclic ring having from 3 to 8 carbon atoms, preferably 5 or 6, optionally substituted.

heterocyclic – represents, in addition to the heteroaryl groups defined below, saturated and unsaturated cyclic organic groups having at least one O, S and/or N atom interrupting a carbocyclic ring structure that consists of one ring or two fused rings, wherein each ring is 3 to 9-membered and may or may not have double bonds that lack delocalized pi electrons, which ring structure has from 2 to 8, preferably from 3 to 6 carbon atoms, e.g., 2- or 3-piperidiny, 2- or 3-piperaziny, 2- or 3-morpholiny, or 2- or 3-thiomorpholiny;

halogen – represents fluorine, chlorine, bromine and iodine;

heteroaryl – represents a cyclic organic group having at least one O, S and/or N atom interrupting a carbocyclic ring structure and having a sufficient number of delocalized pi electrons to provide aromatic character, with the aromatic heterocyclic group having from 2 to 14, preferably 4 or 5 carbon atoms, e.g., 2-, 3- or 4-pyridyl, 2- or 3-furyl, 2- or 3-thienyl, 2-, 4- or 5-thiazolyl, 2- or 4-imidazolyl, 2-, 4- or 5-pyrimidinyl, 2-pyrazinyl, or 3- or 4-pyridazinyl, etc. Preferred heteroaryl groups are 2-, 3- and 4-pyridyl; Such heteroaryl groups may also be optionally substituted.

Also included in the invention are tautomers, enantiomers and other optical isomers of compounds of Formula I or Formula II, as well as pharmaceutically acceptable salts and solvates thereof.

A further feature of the invention is pharmaceutical compositions containing as active ingredient a compound of Formula I or Formula II (or its salt, solvate or isomers) together with a pharmaceutically acceptable carrier or excipient.

The invention also provides methods for preparing compounds of Formula I and Formula II, as well as methods for treating diseases such as, for example, HCV and related disorders. The methods for treating comprise administering to a patient suffering from said disease or diseases a therapeutically effective amount of a compound of Formula I or Formula II, or pharmaceutical compositions comprising a compound of Formula I or Formula II.

Also disclosed is the use of a compound of Formula I and Formula II for the manufacture of a medicament for treating HCV and related disorders.

Detailed description of preferred embodiments

5 In one embodiment, the present invention discloses compounds of Formula I and Formula II as inhibitors of HCV protease, especially the HCV NS3/NS4a serine protease, or a pharmaceutically acceptable derivative thereof, where the various definitions are given above.

10 Representative compounds of the invention which exhibit excellent HCV protease inhibitory activity are listed below along with their activity (ranges of K_i^* values in nanomolar, nM). The Example numbers refer to the numbers for the various structures in the EXAMPLES section found in the later parts of this application.

Table 1: HCV protease continuous assay results

| Example Number | K_i^* (nM) |
|----------------|--------------|
| 1 | c |
| 2 | c |
| 3 | c |
| 4 | b |
| 5 | b |
| 6 | c |
| 7 | c |
| 8 | c |
| 9 | c |
| 10 | c |
| 11 | c |
| 12 | b |

| | |
|----|---|
| 13 | c |
| 14 | c |
| 15 | c |
| 16 | c |
| 17 | c |
| 18 | a |
| 19 | c |
| 20 | c |
| 21 | c |
| 22 | c |
| 23 | a |
| 24 | c |
| 25 | c |
| 26 | c |
| 27 | c |
| 28 | a |
| 29 | c |
| 30 | b |
| 31 | c |
| 32 | c |
| 33 | b |
| 34 | b |
| 35 | c |
| 36 | b |
| 37 | c |
| 38 | c |
| 39 | c |
| 40 | c |
| 41 | c |
| 42 | c |


00000077 0710001





| | |
|----|---|
| 43 | c |
| 44 | c |
| 45 | c |
| 46 | c |
| 47 | c |
| 48 | b |
| 49 | a |
| 50 | b |
| 51 | a |
| 52 | a |
| 53 | a |
| 54 | a |
| 55 | a |
| 56 | a |
| 57 | a |


HCV continuous assay K_i^* range:

a = 1-1000 nM; b = 1001 nM- 50000 nM; and c >50001 nM.


- Some of the types of the inventive compounds and methods of synthesizing the various types of the inventive compounds of Formula I and Formula II are listed below, then schematically described, followed by the illustrative Examples.









6




7




8




| Compound | X ² |
|----------|-------------------------------------------------------|
| 9 | Boc-Val- (isomer 1) |
| 10 | Boc-Val- (isomer 2) |
| 11 | Ac-Glu(O ^t Bu)-Glu(O ^t Bu)-Val- |
| 12 | Ac-Glu-Glu-Val- |
| 13 | Boc |
| 14 | H |




| Compound | X ³ |
|-----------|-------------------------------------------------------|
| 15 | Boc-Val- |
| 16 | Val- |
| 17 | Ac-Glu(O ^t Bu)-Glu(O ^t Bu)-Val- |
| 18 | Ac-Glu-Glu-Val- |



| Compound | X ⁴ | Y ¹ |
|-----------|-------------------------------------------------------|----------------|
| 19 | Boc- | allyl |
| 20 | H | allyl |
| 21 | Boc-Val- | allyl |
| 22 | Ac-Glu(O ^t Bu)-Glu(O ^t Bu)-Val- | allyl |
| 23 | Ac-Glu-Glu-Val- | allyl |
| 24 | Boc-Val- | H |
| 25 | Ac-Glu(O ^t Bu)-Glu(O ^t Bu)-Val- | H |




| Compound | X ⁵ | Y ² | P ₃ |
|----------|-------------------------------------------------------|----------------|----------------|
| 26 | Boc-Val- | allyl | iPr |
| 27 | Ac-Glu(O ^t Bu)-Glu(O ^t Bu)-Val- | allyl | iPr |
| 28 | Ac-Glu-Glu-Val- | allyl | iPr |
| 29 | Boc- | benzyl | Chx |



| Compound | X ⁶ | Z ¹ | Y ⁵ |
|----------|----------------|----------------|------------------|
| 30 | Boc- | 3,4-dichloro | -NH ₂ |
| 31 | Boc- | 4-Bromo | -OBn |
| 32 | Boc- | 3-Bromo | -OBn |
| 33 | Boc- | 4-Chloro | -OBn |
| 34 | Boc- | 3-Chloro | -OBn |
| 35 | iBoc- | 3,4-dichloro | -NH ₂ |
| 36 | iBoc- | 4-Bromo | -OBn |
| 37 | iBoc- | 3-Bromo | -OBn |
| 38 | iBoc- | 4-Chloro | -OBn |
| 39 | iBoc- | 3-Chloro | -OBn |
| 40 | Boc- | 4-Bromo | -OH |
| 41 | Boc- | 3-Bromo | -OH |
| 42 | Boc- | 4-Chloro | -OH |
| 43 | Boc- | 3-Chloro | -OH |
| 44 | iBoc- | 4-Bromo | -OH |
| 45 | iBoc- | 3-Bromo | -OH |
| 46 | iBoc- | 4-Chloro | -OH |

| | | | |
|----|-------|----------|-----|
| 47 | iBoc- | 3-Chloro | -OH |
|----|-------|----------|-----|



| Compound | X ^Z | Z ² |
|----------|----------------|-----------------------|
| 48 | iBoc- | H |
| 49 | Ac-Val- | H |
| 50 | iBoc- | 3,4-dimethyl |
| 51 | Ac-Val- | 3,4-dimethyl |
| 52 | iBoc- | 3-methyl |
| 53 | Ac-Val- | 3-methyl |
| 54 | Ac-Val- | 4-methyl |
| 55 | Ac-Val- | 3-methoxy |
| 56 | Ac-Val- | 4-methoxy |
| 57 | Ac-Val- | 4 ^L -butyl |

- 5 Depending upon their structure, the compounds of the invention may form pharmaceutically acceptable salts with organic or inorganic acids, or organic or inorganic bases. Examples of suitable acids for such salt formation are hydrochloric, sulfuric, phosphoric, acetic, citric, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and carboxylic
- 10 acids well known to those skilled in the art. For formation of salts with bases, suitable bases are, for example, NaOH, KOH, NH₄OH, tetraalkylammonium hydroxide, and the like.

In another embodiment, this invention provides pharmaceutical compositions comprising the inventive peptides as an active ingredient. The

pharmaceutical compositions generally additionally comprise a pharmaceutically acceptable carrier diluent, excipient or carrier (collectively referred to herein as carrier materials). Because of their HCV inhibitory activity, such pharmaceutical compositions possess utility in treating hepatitis C and related disorders.

5 In yet another embodiment, the present invention discloses methods for preparing pharmaceutical compositions comprising the inventive compounds as an active ingredient. In the pharmaceutical compositions and methods of the present invention, the active ingredients will typically be administered in admixture with suitable carrier materials suitably selected with respect to the intended form of administration, i.e. oral tablets, capsules (either solid-filled, semi-solid filled or 10 liquid filled), powders for constitution, oral gels, elixirs, dispersible granules, syrups, suspensions, and the like, and consistent with conventional pharmaceutical practices. For example, for oral administration in the form of tablets or capsules, the active drug component may be combined with any oral 15 non-toxic pharmaceutically acceptable inert carrier, such as lactose, starch, sucrose, cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, talc, mannitol, ethyl alcohol (liquid forms) and the like. Moreover, when desired or needed, suitable binders, lubricants, disintegrating agents and coloring agents may also be incorporated in the mixture. Powders and tablets may be comprised 20 of from about 5 to about 95 percent inventive composition. Suitable binders include starch, gelatin, natural sugars, corn sweeteners, natural and synthetic gums such as acacia, sodium alginate, carboxymethylcellulose, polyethylene glycol and waxes. Among the lubricants there may be mentioned for use in these dosage forms, boric acid, sodium benzoate, sodium acetate, sodium chloride, and 25 the like. Disintegrants include starch, methylcellulose, guar gum and the like. Sweetening and flavoring agents and preservatives may also be included where appropriate. Some of the terms noted above, namely disintegrants, diluents, lubricants, binders and the like, are discussed in more detail below.

30 Additionally, the compositions of the present invention may be formulated in sustained release form to provide the rate controlled release of any one or more of the components or active ingredients to optimize the therapeutic effects, i.e.

HCV inhibitory activity and the like. Suitable dosage forms for sustained release include layered tablets containing layers of varying disintegration rates or controlled release polymeric matrices impregnated with the active components and shaped in tablet form or capsules containing such impregnated or encapsulated porous polymeric matrices.

Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injections or addition of sweeteners and pacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal administration.

Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier such as inert compressed gas, e.g. nitrogen.

For preparing suppositories, a low melting wax such as a mixture of fatty acid glycerides such as cocoa butter is first melted, and the active ingredient is dispersed homogeneously therein by stirring or similar mixing. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool and thereby solidify.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

The compounds of the invention may also be deliverable transdermally. The transdermal compositions may take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type as are conventional in the art for this purpose.

Preferably the compound is administered orally, intravenously or subcutaneously.

Preferably, the pharmaceutical preparation is in a unit dosage form. In such form, the preparation is subdivided into suitably sized unit doses containing

appropriate quantities of the active components, e.g., an effective amount to achieve the desired purpose.

The quantity of the inventive active composition in a unit dose of preparation may be generally varied or adjusted from about 1.0 milligram to about 1,000 milligrams, preferably from about 1.0 to about 950 milligrams, more preferably from about 1.0 to about 500 milligrams, and typically from about 1 to about 250 milligrams, according to the particular application. The actual dosage employed may be varied depending upon the patient's age, sex, weight and severity of the condition being treated. Such techniques are well known to those skilled in the art.

Generally, the human oral dosage form containing the active ingredients can be administered 1 or 2 times per day. The amount and frequency of the administration will be regulated according to the judgment of the attending clinician. A generally recommended daily dosage regimen for oral administration may range from about 1.0 milligram to about 1,000 milligrams per day, in single or divided doses.

Some useful terms are described below:

Capsule - refers to a special container or enclosure made of methyl cellulose, polyvinyl alcohols, or denatured gelatins or starch for holding or containing compositions comprising the active ingredients. Hard shell capsules are typically made of blends of relatively high gel strength bone and pork skin gelatins. The capsule itself may contain small amounts of dyes, opaquing agents, plasticizers and preservatives.

Tablet - refers to a compressed or molded solid dosage form containing the active ingredients with suitable diluents. The tablet can be prepared by compression of mixtures or granulations obtained by wet granulation, dry granulation or by compaction.

Oral gel - refers to the active ingredients dispersed or solubilized in a hydrophilic semi-solid matrix.

Powder for constitution refers to powder blends containing the active ingredients and suitable diluents which can be suspended in water or juices.

Diluent - refers to substances that usually make up the major portion of the composition or dosage form. Suitable diluents include sugars such as lactose, sucrose, mannitol and sorbitol; starches derived from wheat, corn, rice and potato; and celluloses such as microcrystalline cellulose. The amount of diluent in the

composition can range from about 10 to about 90% by weight of the total composition, preferably from about 25 to about 75%, more preferably from about 30 to about 60% by weight, even more preferably from about 12 to about 60%.

Disintegrant - refers to materials added to the composition to help it break apart (disintegrate) and release the medicaments. Suitable disintegrants include starches; "cold water soluble" modified starches such as sodium carboxymethyl starch; natural and synthetic gums such as locust bean, karaya, guar, tragacanth and agar; cellulose derivatives such as methylcellulose and sodium carboxymethylcellulose; microcrystalline celluloses and cross-linked microcrystalline celluloses such as sodium croscarmellose; alginates such as alginic acid and sodium alginate; clays such as bentonites; and effervescent mixtures. The amount of disintegrant in the composition can range from about 2 to about 15% by weight of the composition, more preferably from about 4 to about 10% by weight.

Binder - refers to substances that bind or "glue" powders together and make them cohesive by forming granules, thus serving as the "adhesive" in the formulation. Binders add cohesive strength already available in the diluent or bulking agent. Suitable binders include sugars such as sucrose; starches derived from wheat, corn rice and potato; natural gums such as acacia, gelatin and tragacanth; derivatives of seaweed such as alginic acid, sodium alginate and ammonium calcium alginate; cellulosic materials such as methylcellulose and sodium carboxymethylcellulose and hydroxypropylmethylcellulose; polyvinylpyrrolidone; and inorganics such as magnesium aluminum silicate. The amount of binder in the composition can range from about 2 to about 20% by weight of the composition, more preferably from about 3 to about 10% by weight, even more preferably from about 3 to about 6% by weight.

Lubricant - refers to a substance added to the dosage form to enable the tablet, granules, etc. after it has been compressed, to release from the mold or die by reducing friction or wear. Suitable lubricants include metallic stearates such as magnesium stearate, calcium stearate or potassium stearate; stearic acid; high melting point waxes; and water soluble lubricants such as sodium chloride, sodium benzoate, sodium acetate, sodium oleate, polyethylene glycols and D-leucine. Lubricants are usually added at the very last step before compression, since they must be present on the surfaces of the granules and in between them and the parts of the tablet press. The amount of lubricant in the composition can range from about 0.2 to about 5% by weight of the composition, preferably from about 0.5 to about 2%, more preferably from about 0.3 to about 1.5% by weight.

Glident - material that prevents caking and improve the flow characteristics of granulations, so that flow is smooth and uniform. Suitable glidents include silicon dioxide and talc. The amount of glident in the composition can range from about 0.1% to about 5% by weight of the total composition, preferably from about 0.5 to about 2% by weight.

Coloring agents - excipients that provide coloration to the composition or the dosage form. Such excipients can include food grade dyes and food grade dyes adsorbed onto a suitable adsorbent such as clay or aluminum oxide. The amount of the coloring agent can vary from about 0.1 to about 5% by weight of the composition, preferably from about 0.1 to about 1%.

Bioavailability - refers to the rate and extent to which the active drug ingredient or therapeutic moiety is absorbed into the systemic circulation from an administered dosage form as compared to a standard or control.

Conventional methods for preparing tablets are known. Such methods include dry methods such as direct compression and compression of granulation produced by compaction, or wet methods or other special procedures. Conventional methods for making other forms for administration such as, for example, capsules, suppositories and the like are also well known.

Another embodiment of the invention discloses the use of the pharmaceutical compositions disclosed above for treatment of diseases such as,

for example, hepatitis C and the like. The method comprises administering a therapeutically effective amount of the inventive pharmaceutical composition to a patient having such a disease or diseases and in need of such a treatment.

In yet another embodiment, the compounds of the invention may be used for the treatment of HCV in humans in monotherapy mode or in a combination therapy mode such as, for example, in combination with antiviral agents such as, for example, ribavirin and/or interferon such as, for example, α -interferon and the like.

As stated earlier, the invention includes tautomers, rotamers, enantiomers and other stereoisomers of the compounds also. Thus, as one skilled in the art appreciates, some of the inventive compounds may exist in suitable isomeric forms. Such variations are contemplated to be within the scope of the invention.

Another embodiment of the invention discloses a method of making the compounds disclosed herein. The compounds may be prepared by several techniques known in the art. Representative illustrative procedures are outlined in the following reaction schemes. It is to be understood that while the following illustrative schemes describe the preparation of a few representative inventive compounds, suitable substitution of any of both the natural and unnatural amino acids will result in the formation of the desired compounds based on such substitution. Such variations are contemplated to be within the scope of the invention.

Abbreviations which are used in the descriptions of the schemes, preparations and the examples that follow are:

THF: Tetrahydrofuran

DMF: *N,N*-Dimethylformamide

EtOAc: Ethyl acetate

AcOH: Acetic acid

HOObt: 3-Hydroxy-1,2,3-benzotriazin-4(3*H*)-one

EDCI: 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride

NMM: *N*-Methylmorpholine

MeOH: Methanol

EtOH: Ethanol

iPrOH: isopropanol

tBuOH: *tert*-Butanol

Et₂O: Diethyl ether

5 DME: Ethyleneglycol dimethylether

Boc: *tert*-Butyloxycarbonyl

iBoc: isobutyloxycarbonyl

Cbz: Benzyloxycarbonyl

Fmoc: 9-Fluorenylmethoxycarbonyl

10 DMSO: Dimethyl sulfoxide

TFA: Trifluoroacetic acid

HOBt: Hydroxybenzotriazole

Hünigs base: Diisopropylethyl amine (DIPEA)

BOP: Benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate

15 HATU: O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium
hexafluorophosphate

TsOH: *p*-Toluenesulfonic acid

DMAP: 4-Dimethyl aminopyridine

DCC: Dicyclohexylcarbodiimide

20 MeCN: acetonitrile

Me: Methyl

Bn: Benzyl

Pr: Propyl


iPr: isopropyl

25 Chx: Cyclohexyl


^tBu: *tert*-Butyl

Ac: Acetyl

Ph: Phenyl


Preparation of Intermediates:**Intermediate A:****Step 1:**

To a stirred solution of 1-nitrobutane (16.5 g, 0.16 mol) and glyoxylic acid in H₂O (28.1 g, 0.305 mol) and MeOH (122 mL) at 0°C-5°C, was added dropwise triethyl amine (93 mL, 0.667 mol) over 2 hrs. The solution was warmed to room temperature, stirred overnight and concentrated to dryness to give an oil. The oil was then dissolved in H₂O and acidified to pH =1 with 10% HCl, followed by extraction with EtOAc. The combined organic solution was washed with brine, dried over Na₂SO₄, filtered and concentrated to dryness to give the product **ii** (28.1 g, 99% yield).

Step 2:


To a stirred solution of starting material **ii** (240 g, 1.35 mol) in acetic acid (1.25 L) was added 10% Pd/C (37 g). The resulting solution was hydrogenated at 59 psi for 3 hrs and then at 60 psi overnight. The acetic acid was then evaporated and azeotroped 3 times with toluene, then triturated with MeOH and ether. The solution was then filtered and azeotroped twice with toluene to give an off white solid (131 g, 0.891 mol, 66%).

Step 3:




To a stirred solution of the amino acid **iii** (2.0 g, 13.6 mmol) in dioxane (10 mL) and H₂O (5 mL) at 0°C, was added 1N NaOH solution (4.3 mL, 14.0 mmol). The resulting solution was stirred for 10 minutes, followed by addition of di-*t*-butyldicarbonate (0.110 g, 14.0 mmol) and stirred at 0°C for 15 minutes. The solution was then warmed to room temperature, stirred for 45 minutes and kept at refrigerator overnight and concentrated to dryness to give a crude material. To the solution of this crude material in EtOAc (100 mL) and ice, was added KHSO₄ (3.36 g) and H₂O (32 mL) and stirred for 4-6 minutes. The organic layer was then separated and the aqueous layer was extracted twice with EtOAc and the combined organic layer was washed with water, brine, dried over Na₂SO₄, filtered and concentrated to dryness to give the product as a clear gum (3.0 g, 89% yield).


Step 4:




To a stirred solution of starting material **iv** (6.0 g, 24.3 mmol) in CH₂Cl₂ (120 mL) was added BOP reagent (12.9 g, 29.1 mmol), allyl amine (2.2 mL, 29.1 mmol) and Et₃N (12.0 mL, 85.0 mmol). The resulting solution was stirred at room temperature overnight, then washed with 10% aq. citric acid, satd. NaHCO₃, and brine. The organic layer was dried over Na₂SO₄, concentrated to dryness and ran through column to give the product **v** (5.5 g, 80% yield).

Step 5:

The solution of starting material **v** (2.1 g, 7.3 mmol) in 4M HCl/Dioxane (25 mL) was stirred at room temperature for 1 minute, then concentrated to dryness and dried overnight to give a crude product **A** (2.1 g, 100% yield).

Intermediate B:Step 1:

To a stirred solution of compound **vi** (4.83 g, 19.71 mmol) in dichloromethane (100 mL) at -20°C was added HOObt (3.38 g, 20.70 mmol), *N*-methylmorpholine (8.66 mL, 78.84 mmol) and EDCI (4.90 g, 25.62 mmol). The reaction mixture was stirred for 20 minutes, followed by the addition of **vii** (5.0 g, 19.71 mmol). The resulting solution was stirred at -20° to -10°C for 45 min, then kept in the freezer overnight. The reaction mixture was washed twice with saturated NaHCO₃, 1N HCl, and brine, dried over Na₂SO₄, filtered and concentrated to dryness to give the product **viii** (8.2 g, 94% yield) which was sufficiently pure for further studies. LRMS *m/z* MH⁺ = 445.1.


Step 2:

To a cold (0°C) solution of **viii** (2.63 g, 5.92 mmol) in MeOH (30 mL) was added aqueous 1M LiOH solution (5.92 mL, 5.92 mmol). The reaction mixture was warmed to ambient temperature over 3.5 hrs when the starting material was completely consumed (by TLC). The solvent was removed and the residue was

partitioned between EtOAc and aqueous KHSO_4 (pH 3-4). The organic layer was separated, washed with brine, dried (MgSO_4) and concentrated to afford 2.37 g (93%) of **B**, which was used in the subsequent step without purification. LRMS m/z $\text{MH}^+ = 431.3$.


5 Intermediate C:

Step 1:



The starting material **B** was coupled with commercially available *p*-TsOH.H-Val-OBn using the procedure described for intermediate B, Step 1 to provide **ix** in 94% yield. The crude product was sufficiently pure for further studies. LRMS m/z $\text{MH}^+ = 620.1$.


Step 2:



A solution of the starting material **ix** (1.35 g, 2.18 mmol) in absolute ethanol (20 mL) was stirred at ambient temperature under hydrogen atmosphere in the presence of Pd-C (30 mg, 10%) overnight. The mixture was filtered through a celite pad and the resulting solution was concentrated in vacuo to give the desired product **C** in quantitative yield. LRMS m/z $\text{MH}^+ = 530.1$.

20 Intermediate D:


Step 1:



The desired compound **iv** was obtained by the procedure described for Intermediate A, Step 4 using *p*-TsOH.H-Gly-Oallyl as the coupling partner.

Purification by column chromatography using 50/50 EtOAc/hexanes afforded pure **iv** in 85% yield. LRMS m/z MH^+ = 345.2.


Step 2:




The desired intermediate **D** was obtained by the method described for Intermediate A, Step 5 (reaction time = 1 hr). The crude material was used without further purification.

Intermediate E:


Step 1:



To a stirred solution of starting material (3.00 g, 12.0 mmol) in DMF (15 mL) and CH_2Cl_2 (15 mL) at -20°C was added HOObt (1.97 g, 12.0 mmol), *N*-methyl morpholine (4.0 mL, 36.0 mmol) and EDCI (2.79 g, 14.5 mmol) and stirred for 10 minutes, followed by addition of $\text{HCl}\cdot\text{H}_2\text{N}\text{-Gly-OBn}$ (2.56 g, 13.0 mmol). The resulting solution was stirred at -20°C for 2 hrs, then kept at refrigerator overnight and concentrated to dryness, followed by dilution with EtOAc (150 mL). The EtOAc solution was then washed twice with saturated NaHCO_3 , H_2O , 5% H_3PO_4 , brine, dried over Na_2SO_4 , filtered and concentrated to dryness to give the product (4.5 g, 94%). LRMS m/z MH^+ = 395.1.


Step 2:

The desired intermediate **E** was prepared from compound **xi** according to the procedures described in Step 5 for intermediate **A**. It was used without further purification.


Intermediate F:**Step 1:**

The solution of starting material **xi** (7.00 g, 17.8 mmol) in absolute ethanol (300 mL) was stirred at room temperature under a hydrogen atmosphere in the presence of Pd-C (300 mg, 10%). The reaction progress was monitored by tlc. After 2h, the mixture was filtered through a celite pad and the resulting solution was concentrated in vacuo to give the product **xii** (5.40 g, quantitative). LRMS m/z


$\text{MH}^+ = 305.1$.

Step 2:

The desired product **xiv** was obtained by the procedure described for Intermediate B, Step 1 using commercially available **xiii** as the coupling partner. The crude material was sufficiently pure for further studies.

Step 3:

The desired product **F** was obtained by the procedure described for Intermediate A, Step 5. The crude material was used for further studies.

Intermediate G:**Step 1:**


The desired compound was synthesized by the procedure described for Intermediate E, Step 1 using commercially available **xv** and ammonium chloride. The crude residue was purified by column chromatography using 2.5/97.5 MeOH/dichloromethane to provide **xvi** in 44% yield. LRMS m/z MH^+ = 251.1.

5 Step 2:




The desired compound **xvii** was obtained in quantitative yield using the procedure described for Intermediate A, Step 5. The crude material was used for further studies.


10 Step 3:



15 The desired product **xviii** was obtained by the coupling procedure described for Intermediate E, Step 1 using **xii** and **xvii**. The crude material was sufficiently pure for further studies. LRMS m/z MH^+ = 437.2.


Step 4:

The desired intermediate **G** was obtained from **xviii** using the procedure described for Intermediate A, Step 5 in quantitative yield. The crude material was used for further studies without purification.

Intermediate H:**Step 1:**

The starting material **iii** (21.5 g, 122 mmol) was dissolved in a solution of sodium carbonate (40.76 g) in water (625 mL). Dioxane (200mL) was added. Fmoc-OSu (Fmoc-succinimide, 40.15 g, 121 mmol) in dioxane (400 mL) was added and the reaction mixture was stirred at ambient temperature overnight. Water (1000 mL) was added and washed with ether (6 times) to remove any unreacted Fmoc-OSu. The aqueous layer was acidified (conc. HCl) and extracted with EtOAc. The organic layer was washed with 1N HCl, dried (Na₂SO₄), filtered and concentrated to afford **xix** in quantitative yield.

Step 2:




Commercially available Fmoc-Phg-SASRIN resin (200-400 mesh, 6 g, 0.39 mmol/g) was washed with DMF (2 x 90mL). It was treated with 20% piperidine in DMF (v/v, 100 mL) for 30 min, followed by washing with DMF (4 x 90 mL). DMF (90 mL) was added to the resin followed by Fmoc-Gly-OH (2.09 g, 7.02 mmol), HATU (2.67 g, 7.02 mmol) and DIPEA (2.45 mL, 14.04 mmol). After shaking at ambient temperature overnight, the resin was washed with DMF (4 x 90 mL) to afford resin-bound compound **xxi**.


The above described sequence of Fmoc deprotection and coupling (with **xix** instead of Fmoc-Gly-OH, reaction time = 4hrs) was repeated to afford resin-bound compound **xxii**. The required intermediate **H** was obtained by treating **xxii** with 20% v/v piperidine in DMF as explained before. This material was stored in the refrigerator prior to use.

Examples

Example 1: Preparation of Compound of Formula 1:




Step A:




A solution of compound **1a** (10.0 g, 37.8 mmol) in pyridine (18 mL), t-BuOH (14 mL) and CHCl₃ (25 mL) was cooled at -15°C for 10 minutes, followed by dropwise addition of phosphorous oxychloride (4 mL). The resulting solution was stirred at -15°C for 30 minutes, then warmed to room temperature for 4 hrs. The solution was then washed with H₂O, 1% cold HCl, saturated NaHCO₃, and brine. The combined organic layer was dried over Na₂SO₄, concentrated to dryness and purified by column chromatography on silica gel, eluting with 35% EtOAc/65% CH₂Cl₂ to give compound **1b** (11.0 g, 91% yield).

Step B:




To a stirred solution of compound **1b** (2.45 g, 15.2 mmol) in DME (35 mL) was added K_2CO_3 (2.1 g, 15.2 mmol), EtO_2C-CH_2Br (3.4 mL, 30.7 mmol) and a small amount of tetra-*n*-butylammonium iodide. The resulting solution was stirred at room temperature overnight and concentrated to dryness, followed by extraction with EtOAc-brine. The combined organic layer was dried over Na_2SO_4 , then concentrated to dryness and purified by column chromatography, eluting with 25% EtOAc/75% CH_2Cl_2 to give compound **1c** (2.06 g, 66% yield).

Step C:




To a stirred solution of starting material **1c** (1.0 g, 2.46 mmol) was added TFA (4.0 mL, 2.46 mmol). The resulting solution was stirred at room temperature for 1 hr, concentrated to dryness and dried overnight to give compound **1d** (0.88 g, 100% yield).

Step D:





To a stirred solution of starting material **1d** (0.3 g, 0.857 mmol) in CH_2Cl_2 (15 mL) was added **A** (0.19 g, 0.857 mmol), BOP reagent (0.42 g, 0.94 mmol) and Et_3N (0.42 mL, 3.0 mmol). The resulting solution was stirred at room temperature overnight, then washed with 10% aqueous citric acid, saturated NaHCO_3 , and brine. The combined organic layer was dried over Na_2SO_4 , concentrated to dryness and then ran through column to give compound **1e** (0.15 g, 34% yield).


Step E:



To a stirred cooling solution of compound **1e** (0.1 g, 0.193 mmol) in CH_2Cl_2 (10 mL) at 0°C , was added Dess-Martin's oxidant (0.123 g, 0.289 mmol). The resulting solution was stirred at room temperature for 1 hour, then poured into aq. $\text{Na}_2\text{S}_2\text{O}_3$ (15 mL), satd. NaHCO_3 (15 mL) and CH_2Cl_2 (15 mL) with ice and stirred at room temperature for half an hour. The solution was then concentrated to dryness and purified by prep TLC using 5% MeOH/ 95% CH_2Cl_2 to give the final product, compound **1** (90 mg, 91% yield). HRMS (FAB) Calcd for $\text{C}_{25}\text{H}_{33}\text{N}_4\text{O}_8$: 517.2298 ($\text{M}+\text{H}$)⁺. Found: 517.2307.


Example 2: Preparation of Compound of Formula 2:**2****Step A:**

The desired product **2a** was prepared in 60% isolated yield, as described above for Example 1, Step B, using ethyl-4-bromobutyrate as the alkylating agent.


Step B:

The desired product **2b** was prepared in quantitative yield, as described above for Example 1, Step C.


Step C:




To a stirred solution of starting material **2b** (0.174 g, 0.46 mmol) in CH₂Cl₂ (5 mL) was added **A** (0.113 g, 0.51 mmol), EDCI (0.106 g, 0.552 mmol), HOBt (0.085g, 0.552 mmol) and NMM (0.177 mL, 1.61 mmol). The resulting solution was stirred at -20°C for 48 hrs. The solution was then washed with satd. NaHCO₃, 10% aq. citric acid and brine. The combined organic layer was dried over Na₂SO₄ and purified by column chromatography using 5/95 MeOH/CH₂Cl₂ to give product **2c** (0.082 g, 71% yield, MH⁺ = 547.3).


Step D:

The desired product **2** was prepared in 78% isolated yield, as described above for Example 1, Step E. HRMS (FAB) Calcd for $C_{27}H_{37}N_4O_8$: 545.2611 (M+H)⁺. Found: 545.2607.


Example 3: Preparation of Compound of Formula 3:

Step A:


- 5 To the starting material **1a** (10.50 g, 39.74 mmol) in benzene (100 mL) was added benzyl alcohol (10.30 mL, 99.35 mmol) and *p*-toluenesulfonic acid (1.52 g, 7.95 mmol). The reaction mixture was refluxed overnight. It was then cooled to room temperature and washed with saturated NaHCO₃ solution and brine. The organic layer was separated, dried (Na₂SO₄) and concentrated. The residue was dissolved in minimum amount of EtOAc and then hexanes was added to this solution. The white solid which crashed out was filtered and dried to give 8.2 g (58%) of pure product **3a**.

Step B:

The desired product **3b** was obtained from **3a** using the procedure described for Example 1, Step B. Purification of the crude material using 10/90 EtOAc/dichloromethane provided **3b** in 74% yield.

Step C:

5 A solution of **3b** (1.29 g, 2.93 mmol) in 30% HBr in AcOH (5 mL) was stirred at ambient temperature for 20 min. It was then poured slowly into a stirred solution of ice/saturated NaHCO₃ and extracted with EtOAc. The organic layer was separated, dried (Na₂SO₄) and concentrated. Purification by flash column chromatography using 30/70 to 50/50 EtOAc/dichloromethane provided 632 mg (70%) of **3c**.


Step D:

15 To a solution of *N*-Boc-valine (424 mg, 1.95 mmol) and *N*-hydroxysuccinimide (230 mg, 2.0 mmol) in THF (8 mL) at 0°-5°C was added DCC (413 mg, 2.0 mmol). The reaction mixture was slowly warmed to ambient temperature overnight. The solid was filtered off and the filtrate was concentrated. The crude *N*-Boc-valine succinimide ester was used further without purification.

20 To a cold (-50°C) solution of **3c** (600 mg, 1.96 mmol) in THF (5 mL) was added potassium tert-butoxide (220 mg, 1.96 mmol) and warmed to -20°C over 20 minutes. To this mixture was added a slurry of the crude *N*-Boc-valine succinimide ester (described above) in THF (4 mL) and warmed to -10°C over 20 minutes. The reaction was quenched with saturated ammonium chloride solution


and extracted with EtOAc. The organic layer was separated, dried (Na_2SO_4) and concentrated. Purification by flash column chromatography using 5/95 to 10/90 EtOAc/dichloromethane provided 543 mg (55%) of **3d** as a mixture of diastereomers.

5 Step E:




To a solution of **3d** (525 mg, 1.04 mmol) in EtOH (8 mL) was added 5% palladium on carbon. The suspension was stirred under hydrogen atmosphere for 2 hrs. The catalyst was filtered off through a pad of celite and the filtrate was concentrated to provide **3e** in quantitative yield.

Step F:




The expected product **3f** was obtained by the protocol described above for Example 2, step C. Purification by flash column chromatography using 5/95 MeOH/dichloromethane provided **3f** in 47% yield as a mixture of diastereomers.



Step G:





The desired product **3** was prepared in 67% isolated yield, as described above for Example 1, Step E.

Example 4: Preparation of Compound of Formula 4:




Step A:

4N HCl in dioxane (5 mL) was added to **3f** (200 mg, 0.343 mmol) and cooled to 0°C. The reaction mixture was slowly warmed to ambient temperature over 3 hrs. It was then concentrated and dried overnight to provide **4a** in quantitative yield, which was used without purification.



Step B:

To a cold (-20°C) solution of **4a** (110 mg, 0.21 mmol) and **C** (111 mg, 0.21 mmol) in anhydrous dichloromethane (4 mL) was added EDCI (48 mg, 0.252 mmol), HOObt (41 mg, 0.252 mmol) and NMM (0.06 mL, 0.525 mmol). After being stirred at this temperature for 2 days the reaction mixture was diluted with dichloromethane and washed with saturated NaHCO_3 , 10% aqueous citric acid, and brine. It was dried over sodium sulfate, filtered and concentrated in vacuo. Flash chromatography (6/94 MeOH/dichloromethane) afforded **4b** (169 mg) in 81% yield.

Step C:



The desired product **4** was obtained by the oxidation of **4b** in 73% isolated yield, as described above for Example 1, Step E.



Example 5: Preparation of Compound of Formula 5:**Step A:**

The desired product **5** was prepared in quantitative yield, as described above for Example 4, Step A.

The following compounds (**6-8**) shown in Table 2 were prepared by essentially the above described procedures using amino acid derivatives from commercial or literature sources.

CCOC(=O)CC1C(=O)N(C(=O)NCC(=O)NCC(=O)NCC=C)C(=O)N1C(=O)C(C)C

| Compound | X ¹ | M+H (HRMS) |
|----------|-------------------------------------------------------|------------|
| 3 | Boc-Val- | 582.3133 |
| 4 | Ac-Glu(O ^t Bu)-Glu(O ^t Bu)-Val- | 993.5470 |
| 5 | Ac-Glu-Glu-Val- | 881.4263 |
| 6 | Ac-Glu(O ^t Bu)- | 709.3776 |
| 7 | Ac-Glu- | 653.3156 |
| 8 | iBoc-Val- | 681.3822 |

Examples 9-14: Preparation of Compounds of Formulae 9-14:**Scheme 1****Table 3**

| Compound | X^2 | M+H (HRMS) |
|-----------|-------------------------------------------------------|------------|
| 9 | Boc-Val- (isomer 1) | 595.3463 |
| 10 | Boc-Val- (isomer 2) | 595.3463 |
| 11 | Ac-Glu(O ^t Bu)-Glu(O ^t Bu)-Val- | 907.5120 |

| | | |
|-----------|-----------------|----------|
| 12 | Ac-Glu-Glu-Val- | 795.3889 |
| 13 | Boc | 496.2779 |
| 14 | H | 396.2247 |

Synthesis of 9a: To a cold (0°C) solution of **3a** (2.0 g, 5.65 mmol) in THF (25 mL) was added NaH (60% dispersion in oil, 264 mg, 6.6 mmol) in two portions. Slowly warmed the reaction mixture to 10°C over 2 hrs. It was quenched with aqueous 10% citric acid solution and brine and extracted into EtOAc. The organic layer was dried over sodium sulfate, filtered and concentrated. The crude material was purified by flash chromatography using 3/97 MeOH/dichloromethane. The material obtained was triturated with minimum amount of ether and filtered. The solid was dried under vacuum overnight to provide pure **9a** (1.31 g) in 66% yield as a white solid.

Synthesis of 9b: The required product **9b** was synthesized using the procedure described for Example 3, Step D in 80% yield, after purification by flash chromatography, as a white foam.

Synthesis of 9c: The required product **9c** was obtained using the procedure described for Example 3, Step E (reaction time – overnight) in quantitative yield as a white foam.

The compounds (**9-14**) shown in Table 3 were prepared by procedures described previously.



Examples 15-18: Preparation of Compounds of Formulae 15-18:**Scheme 2**

Table 4



| Compound | X ³ | M+H (HRMS) |
|-----------|-------------------------------------------------------|------------|
| 15 | Boc-Val- | 685.3900 |
| 16 | Val- | 585.3401 |
| 17 | Ac-Glu(O ^t Bu)-Glu(O ^t Bu)-Val- | 997.5631 |
| 18 | Ac-Glu-Glu-Val- | 885.4359 |

Synthesis of 15c: The starting material **3a** was converted to **15a** and then to **15b** using the literature described procedure (K. Hayashi et. al., *J. Med. Chem.*, **1989**, 32, 289-297). The required product **15c** was obtained from **15b** by the procedure described for Example 3, Step D in 50% isolated yield.

Synthesis of 15d: The required product **15d** was obtained using the procedure described for Example 3, Step E (reaction time – overnight) in quantitative yield.

The compounds (**15-18**) shown in Table 4 were prepared by procedures described previously.

Examples 19-25: Preparation of Compounds of Formulae 19-25:

Scheme 3





Table 5



| Compound | X ⁴ | Y ¹ | M+H (HRMS) |
|-----------|-------------------------------------------------------|----------------|------------|
| 19 | Boc- | allyl | 644.3308 |
| 20 | H | allyl | 544.2762 |
| 21 | Boc-Val- | allyl | 743.3990 |
| 22 | Ac-Glu(O ^t Bu)-Glu(O ^t Bu)-Val- | allyl | 1055.5617 |
| 23 | Ac-Glu-Glu-Val- | allyl | 943.4410 |
| 24 | Boc-Val- | H | 703.3680 |
| 25 | Ac-Glu(O ^t Bu)-Glu(O ^t Bu)-Val- | H | 1015.5353 |

Synthesis of 19a: The required compound **19a** was prepared by the coupling reaction of **15d** and intermediate **D** using the procedure described for Example 4, Step B in 64% isolated yield.

Synthesis of 19: The desired compound **19** was obtained by the oxidation protocol described previously for Example 1, Step E. The crude material was purified by flash column chromatography using 5/1 dichloromethane/EtOAc to provide **19** in 83% yield.

Synthesis of compounds 20-23: Preparation of these derivatives follow essentially previously described procedures.

Synthesis of compounds 24, 25: To a solution of the allyl ester (**21** or **22**, 0.12 mmol) in THF (4 mL) under nitrogen atmosphere was added morpholine (1.2 mmol) and tetrakis(triphenylphosphine)palladium(0) (0.012 mmol). The reaction mixture was kept in the refrigerator (-5° to -10°C) overnight. The solvent was removed in vacuo. The residue was dissolved in dichloromethane and washed

with 5% H_3PO_4 , water and brine. The organic layer was dried over sodium sulfate, filtered and concentrated. Purification by preparative TLC using 5/95 MeOH/dichloromethane provided the required product **24** or **25** (25–40% yield).

Examples 26-29: Preparation of Compounds of Formulae 26-29:

5 **Scheme 4**





Table 6



| Compound | X^5 | Y^2 | P_3 | M+H (LC-MS) |
|-----------|-------------------------------------------------------|---------------------|-------|--------------|
| 26 | Boc-Val- | allyl | iPr | 813.5 |
| 27 | Ac-Glu(O ^t Bu)-Glu(O ^t Bu)-Val- | allyl | iPr | 1125.3 |
| 28 | Ac-Glu-Glu-Val- | allyl | iPr | 1011.2 |
| 29 | Boc- | benzyl ^a | Chx | 702.1 (-100) |

a. Prepared using intermediate **E** as the coupling partner with **26d**.

Synthesis of 26a: The required material **26a** was obtained from **3a** and commercially available 3,4-dichlorobenzyl bromide using the procedure described for the synthesis of **15c** (K. Hayashi et. al., *J. Med. Chem.*, **1989**, 32, 289-297) in 83% yield after purification. HRMS (FAB) Calcd for $C_{26}H_{23}N_2O_5Cl_2$: 513.0984 (M+H)⁺. Found: 513.0991.

Synthesis of **26b** and **26c/d** follow previously described procedures.

Synthesis of 26e: To a solution of **26c** (1.25 g, 2.16 mmol) in EtOH (10 mL) was added 1M aqueous LiOH (3.02 mL, 3.02 mmol) and stirred at ambient temperature for 2 hrs. The reaction mixture was quenched with 10% aqueous citric acid solution (acidic by pH paper) and extracted with chloroform. The organic layer was dried (Na₂SO₄), filtered and concentrated to provide **26e**. The crude material was used further without any purification. (**26f** was prepared by a similar procedure from **26d**).


Synthesis of compounds **26-29** shown above (Table 6) follow essentially previously described procedures.

Examples 30-47: Preparation of Compounds of Formulae 30-47:

General procedure for the N-benylation of imidazolidinone **3a**: The following

- 5 imidazolidinone derivatives **I-1** to **I-10** (Table 7) were obtained by the N-alkylation of **3a** and commercially available substituted benzyl bromide or chloride using the conditions described in literature (K. Hayashi et. al., *J. Med. Chem.*, **1989**, 32, 289-297). In most cases the crude material after alkylation was sufficiently pure for further manipulation and were obtained in almost quantitative yield.

Table 7



| Compound | Z ¹ | HRMS (Calcd / found) |
|----------|-----------------------|-------------------------|
| I-1 | 4-bromo | 523.0869 / 523.0865 |
| I-2 | 3-bromo | 523.0869 / 523.0878 |
| I-3 | 4-chloro | 479.1374 / 479.1370 |
| I-4 | 3-chloro | 479.1374 / 479.1374 |
| I-5 | 3,4-dimethyl | 473.2076 / 473.2088 |
| I-6 | 3-methyl | 459.1920 / 459.1908 |
| I-7 | 4-methyl | 459.1920 / 459.1917 |
| I-8 | 3-methoxy | 475.1869 / 475.1859 |
| I-9 | 4-methoxy | 475.1869 / 475.1873 |
| I-10 | 4- ^t butyl | 501.2 ^a |

a. LRMS data.

Scheme 5





Table 8



| Compound | X ⁶ | Z ¹ | Y ⁵ | M+H (LC-MS) |
|----------|----------------|----------------|------------------|-------------|
| 30 | Boc- | 3,4-dichloro | -NH ₂ | 844.2 |
| 31 | Boc- | 4-Bromo | -OBn | 947.3 |
| 32 | Boc- | 3-Bromo | -OBn | 947.3 |
| 33 | Boc- | 4-Chloro | -OBn | 901.3 |
| 34 | Boc- | 3-Chloro | -OBn | 901.2 |
| 35 | iBoc- | 3,4-dichloro | -NH ₂ | 844.2 |
| 36 | iBoc- | 4-Bromo | -OBn | 947.3 |
| 37 | iBoc- | 3-Bromo | -OBn | 947.3 |
| 38 | iBoc- | 4-Chloro | -OBn | 901.3 |
| 39 | iBoc- | 3-Chloro | -OBn | 901.3 |
| 40 | Boc- | 4-Bromo | -OH | 857.2 |
| 41 | Boc- | 3-Bromo | -OH | 857.2 |
| 42 | Boc- | 4-Chloro | -OH | 813.2 |
| 43 | Boc- | 3-Chloro | -OH | 813.2 |
| 44 | iBoc- | 4-Bromo | -OH | 857.2 |
| 45 | iBoc- | 3-Bromo | -OH | 857.2 |
| 46 | iBoc- | 4-Chloro | -OH | 813.2 |
| 47 | iBoc- | 3-Chloro | -OH | 813.2 |


- 5 **Synthesis of 30e-i:** The precursors (**26f**, **I-1** – **I-4**) were prepared by earlier described protocol from appropriate building blocks. Coupling of these with intermediate **F** (Y³ = OBn) or **G** (Y³ = NH₂) independently using the

EDCI/HOObt/NMM procedure described previously provided the desired compounds **30e-i**.

Synthesis of 30-34: The required compounds **30-34** were obtained by the oxidation procedure described for Example 1, Step E. Minimum amount of DMSO was added as cosolvent in cases where the starting material was not completely soluble in dichloromethane. Purification involved flash chromatography using MeOH (1-6%) in dichloromethane as the solvent system.

Synthesis of 35-39: The appropriate starting material (**30-34**, 0.1 mmol) was treated with 4M HCl in dioxane (3-5 mL) at 0°C for 2 hrs. The reaction mixture was concentrated to afford the crude amine hydrochloride. This material was dissolved in THF (5 mL) and cooled to 0°C. To the cold solution was added triethylamine (0.44 mmol) and isobutylchloroformate (0.122 mmol). The reaction mixture was left in the freezer (~ -8°C) overnight. The solvent was evaporated and the residue was dissolved in EtOAc, washed with water, saturated NaHCO₃, saturated NH₄Cl and brine. The organic layer was dried (Na₂SO₄), filtered and concentrated to afford **35-39** which were homogeneous by TLC.

Synthesis of 40-47: To a solution of the appropriate starting material (**31-34**, **36-39**, 0.05 mmol) in THF (3mL) was added a solution of K₂CO₃ (0.06 mmol) in water (3mL). If required, THF (1 mL) was added to obtain a homogeneous solution. t-Butanol (0.1 mL) was added after one hr and the reaction mixture was stirred at ambient temperature till completion (few hrs to 2 days). The reaction was quenched with 10% aqueous citric acid (acidic by pH paper) and extracted into dichloromethane. The organic layer was dried (Na₂SO₄), filtered and concentrated to afford **40-47**.

Examples 48-57: Preparation of Compounds of Formulae 48-57:**Scheme 6**

Synthesis of 49a-j: The precursors **48a-j** were synthesized from appropriate building blocks (**15a, 1-5 – 1-10**) using the procedures described earlier. Removal of the benzyl ester to provide the required compounds **49a-j** were carried out using the procedure described for Example 3, Step E (reaction time = 2-3 hrs) in quantitative yield.

Synthesis of 50a-j: Coupling reaction of **49a** (0.12 mmol) with resin-bound compound **H** (0.039 mmol) was carried out as described for Intermediate **H**, Step 2 using HATU/DIPEA (0.12 mmol/0.24 mmol, reaction time = 4 hrs) in DMF (2 mL). The reaction mixture was washed with DMF (4 x 4 mL), THF (2 x 4 mL) and dichloromethane (2 x 4 mL) to afford resin-bound compound **50a**. Similar procedure was employed for the synthesis of **50b-j**.

Synthesis of 51a-j: Resin-bound compound **50a** (0.039 mmol) was treated with a solution of Dess-Martin's periodinane (0.16 mmol) and t-butanol (0.16 mmol) in dichloromethane (3 mL) at ambient temperature for 4 hrs. The resin was washed with 20% v/v iPrOH in dichloromethane (2 x 4 mL), THF (2 x 4 mL), 50%

v/v aqueous THF (2 x 4 mL), THF (2 x 4 mL), and dichloromethane (4 x 4 mL) to provide resin-bound compound **51a**. Similar procedure was employed for the synthesis of **51b-j**.

- Synthesis of 48-57: To the resin-bound compound **51a** (0.039 mmol) was added 5% v/v TFA in dichloromethane (4 mL). After shaking for 5 minutes, the reaction mixture was filtered into a vial containing AcOH (1 mL). The filtrate was concentrated by vacuum centrifugation to afford the desired compound **48**. Similar procedure was used for the synthesis of **49-57**.

Scheme 7





Table 9



| Compound | X' | Z' | M+H (LC-MS) |
|----------|---------|------------------------|-------------|
| 48 | iBoc- | H | 777.2 |
| 49 | Ac-Val- | H | 818.2 |
| 50 | iBoc- | 3,4-dimethyl | 805.2 |
| 51 | Ac-Val- | 3,4-dimethyl | 846.2 |
| 52 | iBoc- | 3-methyl | 791.2 |
| 53 | Ac-Val- | 3-methyl | 832.2 |
| 54 | Ac-Val- | 4-methyl | 832.2 |
| 55 | Ac-Val- | 3-methoxy | 848.2 |
| 56 | Ac-Val- | 4-methoxy | 848.2 |
| 57 | Ac-Val- | 4- ^t -butyl | 874.2 |

5 **Assay for HCV Protease Inhibitory Activity:**

Spectrophotometric Assay: Spectrophotometric assay for the HCV serine protease was performed on the inventive compounds by following the procedure described by R. Zhang *et al*, *Analytical Biochemistry*, **270** (1999) 268-275, the disclosure of which is incorporated herein by reference. The assay based on the proteolysis of chromogenic ester substrates is suitable for the continuous monitoring of HCV NS3 protease activity. The substrates were derived from the P side of the NS5A-NS5B junction sequence (Ac-DTEDVVX(Nva), where X = A or P) whose C-terminal carboxyl groups were esterified with one of four different chromophoric alcohols (3- or 4-nitrophenol, 7-hydroxy-4-methyl-coumarin, or 4-phenylazophenol). Presented below are the synthesis, characterization and

application of these novel spectrophotometric ester substrates to high throughput screening and detailed kinetic evaluation of HCV NS3 protease inhibitors.

Materials and Methods:

Materials: Chemical reagents for assay related buffers were obtained from

- 5 Sigma Chemical Company (St. Louis, Missouri). Reagents for peptide synthesis were from Aldrich Chemicals, Novabiochem (San Diego, California), Applied Biosystems (Foster City, California) and Perseptive Biosystems (Framingham, Massachusetts). Peptides were synthesized manually or on an automated ABI model 431A synthesizer (from Applied Biosystems). UV/VIS Spectrometer model
- 10 LAMBDA 12 was from Perkin Elmer (Norwalk, Connecticut) and 96-well UV plates were obtained from Corning (Corning, New York). The prewarming block was from USA Scientific (Ocala, Florida) and the 96-well plate vortexer was from Labline Instruments (Melrose Park, Illinois). A Spectramax Plus microtiter plate reader with monochromator was obtained from Molecular Devices (Sunnyvale, California).

Enzyme Preparation: Recombinant heterodimeric HCV NS3/NS4A protease (strain 1a) was prepared by using the procedures published previously (D. L. Sali *et al*, *Biochemistry*, 37 (1998) 3392-3401). Protein concentrations were determined by the Biorad dye method using recombinant HCV protease standards

20 previously quantified by amino acid analysis. Prior to assay initiation, the enzyme storage buffer (50 mM sodium phosphate pH 8.0, 300 mM NaCl, 10% glycerol, 0.05% lauryl maltoside and 10 mM DTT) was exchanged for the assay buffer (25 mM MOPS pH 6.5, 300 mM NaCl, 10% glycerol, 0.05% lauryl maltoside, 5 μ M EDTA and 5 μ M DTT) utilizing a Biorad Bio-Spin P-6 prepacked column.

- 25 Substrate Synthesis and Purification: The synthesis of the substrates was done as reported by R. Zhang *et al*, (*ibid.*) and was initiated by anchoring Fmoc-Nva-OH to 2-chlorotrityl chloride resin using a standard protocol (K. Barlos *et al*, *Int. J. Pept. Protein Res.*, 37 (1991), 513-520). The peptides were subsequently assembled, using Fmoc chemistry, either manually or on an automatic ABI model 431 peptide synthesizer. The N-acetylated and fully protected peptide fragments were cleaved
- 30 from the resin either by 10% acetic acid (HOAc) and 10% trifluoroethanol (TFE) in

dichloromethane (DCM) for 30 min, or by 2% trifluoroacetic acid (TFA) in DCM for 10 min. The combined filtrate and DCM wash was evaporated azeotropically (or repeatedly extracted by aqueous Na_2CO_3 solution) to remove the acid used in cleavage. The DCM phase was dried over Na_2SO_4 and evaporated.

5 The ester substrates were assembled using standard acid-alcohol coupling procedures (K. Holmber *et al*, *Acta Chem. Scand.*, B33 (1979) 410-412). Peptide fragments were dissolved in anhydrous pyridine (30-60 mg/ml) to which 10 molar equivalents of chromophore and a catalytic amount (0.1 eq.) of para-toluenesulfonic acid (pTSA) were added. Dicyclohexylcarbodiimide (DCC, 3 eq.)
10 was added to initiate the coupling reactions. Product formation was monitored by HPLC and found to be complete following 12-72 hour reaction at room temperature. Pyridine solvent was evaporated under vacuum and further removed by azeotropic evaporation with toluene. The peptide ester was deprotected with 95% TFA in DCM for two hours and extracted three times with anhydrous ethyl
15 ether to remove excess chromophore. The deprotected substrate was purified by reversed phase HPLC on a C3 or C8 column with a 30% to 60% acetonitrile gradient (using six column volumes). The overall yield following HPLC purification was approximately 20-30%. The molecular mass was confirmed by electrospray ionization mass spectroscopy. The substrates were stored in dry powder form
20 under desiccation.

Spectra of Substrates and Products: Spectra of substrates and the corresponding chromophore products were obtained in the pH 6.5 assay buffer. Extinction coefficients were determined at the optimal off-peak wavelength in 1-cm cuvettes (340 nm for 3-Np and HMC, 370 nm for PAP and 400 nm for 4-Np) using multiple
25 dilutions. The optimal off-peak wavelength was defined as that wavelength yielding the maximum fractional difference in absorbance between substrate and product (product OD - substrate OD)/substrate OD).

Protease Assay: HCV protease assays were performed at 30°C using a 200 µl reaction mix in a 96-well microtiter plate. Assay buffer conditions (25 mM MOPS
30 pH 6.5, 300 mM NaCl, 10% glycerol, 0.05% lauryl maltoside, 5 µM EDTA and 5 µM DTT) were optimized for the NS3/NS4A heterodimer (D. L. Sali *et al*, *ibid.*)).

Typically, 150 μ l mixtures of buffer, substrate and inhibitor were placed in wells (final concentration of DMSO 4 % v/v) and allowed to preincubate at 30 °C for approximately 3 minutes. Fifty μ l of prewarmed protease (12 nM, 30°C) in assay buffer, was then used to initiate the reaction (final volume 200 μ l). The plates were monitored over the length of the assay (60 minutes) for change in absorbance at the appropriate wavelength (340 nm for 3-Np and HMC, 370 nm for PAP, and 400 nm for 4-Np) using a Spectromax Plus microtiter plate reader equipped with a monochromator (acceptable results can be obtained with plate readers that utilize cutoff filters). Proteolytic cleavage of the ester linkage between the Nva and the chromophore was monitored at the appropriate wavelength against a no enzyme blank as a control for non-enzymatic hydrolysis. The evaluation of substrate kinetic parameters was performed over a 30-fold substrate concentration range (~6-200 μ M). Initial velocities were determined using linear regression and kinetic constants were obtained by fitting the data to the Michaelis-Menten equation using non-linear regression analysis (Mac Curve Fit 1.1, K. Raner). Turnover numbers (k_{cat}) were calculated assuming the enzyme was fully active.

Evaluation of Inhibitors and Inactivators: The inhibition constants (K_i) for the competitive inhibitors Ac-D-(D-Gla)-L-I-(Cha)-C-OH (27), Ac-DTEDVVA(Nva)-OH and Ac-DTEDVVP(Nva)-OH were determined experimentally at fixed concentrations of enzyme and substrate by plotting v_o/v_i vs. inhibitor concentration ($[I]_o$) according to the rearranged Michaelis-Menten equation for competitive inhibition kinetics: $v_o/v_i = 1 + [I]_o / (K_i (1 + [S]_o / K_m))$, where v_o is the uninhibited initial velocity, v_i is the initial velocity in the presence of inhibitor at any given inhibitor concentration ($[I]_o$) and $[S]_o$ is the substrate concentration used. The resulting data were fitted using linear regression and the resulting slope, $1/(K_i(1+[S]_o/K_m))$, was used to calculate the K_i^* value.

The obtained K_i^* values for the various compounds of the present invention are given in the afore-mentioned **Table** wherein the compounds have been arranged in the order of ranges of K_i^* values. From these test results, it would be apparent to the skilled artisan that the compounds of the invention have excellent utility as NS3-serine protease inhibitors.

While the present invention has been described with in conjunction with the specific embodiments set forth above, many alternatives, modifications and other variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present invention.

100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
2233
2234
2235
2236
2237
2238
2239
2240
2241
2242
2243
2244
2245
2246
2247
2248
2249
2250
2251
2252
2253
2254
2255
2256
2257
2258
2259
2260
2261
2262
2263
2264
2265
2266
2267
2268
2269
2270
2271
2272
2273
2274
2275
2276
2277
2278
2279
2280
2281
2282
2283
2284
2285
2286
2287
2288
22